

Sports Therapy
Manual Therapy
Electrotherapy
Manipulation
Rehabilitation
Pre-habilitation
Clinical Pilates
Pre-Season Screening
Exercise Prescription
Massage

Partners

Chris Ward
B.App.Sc. (Physio)

Charlie Schall-
Riaucour
B.Physio (Hons.),
B.App.Sc. (Phys.Ed.),
Post Grad Dip (Exs for
Rehab)

Physiotherapists

Richard Webber
B.App.Sc (Physio)

Sarah Tulloch
B. Health Sc., M.
Physio

Eleanor Dwyer
B. Sc., M. Physio

Daniel Ford
B. Physio

Elsa Gales
BSc. (Physio)

Practice Manager

Anne Mellor
BA (Hons)

POST-OPERATIVE PHYSIOTHERAPY INFORMATION

Shoulder Stabilisation – Mr. Richard Dallalana

Surgery:

Superficially there may only be small puncture marks made by the surgical instruments and these have been closed by steri-strips or a single stitch. Although this may seem only minor, the surgery inside the joint is more traumatic and must be given full opportunity to recover. Your surgeon has repaired the ligaments and cartilage (labrum) in the shoulder that are damaged during dislocation of the joint. Sometimes there is a larger incision over the front of the shoulder to gain access to these structures.

These repaired structures need protection following the surgery in order to heal properly. **You must limit your external rotation of your shoulder to a neutral position for 4 weeks, or until otherwise advised by your surgeon.** If in doubt, keep the arm across your body.

Use of Sling:

The sling may be required to rest and support the shoulder for a period of 4 weeks following surgery, or until your surgeon advises you. During this time you must remove your arm from the sling 3 - 4 times per day to perform the following exercises as demonstrated by your physiotherapist. You may like to use the sling for a little while longer whilst in crowds if your shoulder remains painful as a “warning sign” for others to keep clear!

Exercises: Are to be performed 3 - 4 times per day. The following exercises have been prescribed for you by Mr Dallalana to maintain the movement in the joints of your arm and prevent the shoulder from stiffening too much in the first few weeks. They are to be performed 3-4 times each day and are not to be forced into pain.

1.Elbow, Wrist and Hand Movements – Day 1 onwards

Be sure to keep the elbow, wrist and hand mobile by bending and straightening all joints, keeping your palm facing backwards to protect your shoulder. It is much easier to keep these joints moving now than to let them get stiff. You can also begin gripping a stress ball or squash ball to keep your arm muscles strong and begin activity around the shoulder.

2.Pendular Swings - Day 1 onwards


Stand beside a table/ bench and support yourself firmly with your uninjured arm. Bend forward from the hips to allow your injured arm to hang free. Swing the arm gently forward/ backwards 10 times, side to side 10 times and in circles 10 times. Always keep your palm facing backwards to avoid stressing the shoulder site.



Member

3. Shoulder (scapula) retraction – Day 1 onwards

Pull shoulder blades back gently and pinch them together. Keep shoulders down away from ears. Hold for 5 seconds and repeat 10 times.



4. Assisted shoulder flexion – Day 1/ 2 onwards

Lying on your back, cradle your injured arm in your uninjured arm. Raise both together in front using the uninjured arm to assist the injured arm. Move only to the point of pain. Make sure that you don't let your shoulder "hitch" upwards when elevating the arm. Repeat this 10 times



★ Do not elevate your operated arm past 90° unless directed by your surgeon and/or physiotherapist.

Bandage/ Dressings

The wounds will be covered by waterproof dressings prior to discharge from hospital. You need to leave these on until your review with your surgeon. Once the wound has healed sufficiently (7-10 days), you are able to shower without the waterproof dressings.

Further Rehabilitation/ Return to Normal Activity

It is advisable for you to attend a physiotherapist for a progressive regime of exercises to mobilise and strengthen your shoulder for return to normal function and/ or sport. The shoulder is a complicated joint that requires intricate muscle function to perform normally, especially for the demands of overhead sporting or work activities, and physiotherapy can help guide your return to these activities. Your surgeon will advise you at the time of follow up review when you should commence further physiotherapy.

Return to full sporting activities is usually between 3-6 months, depending upon the nature of your chosen sport and how well your strength and movements have progressed. Return to work is very dependant on your occupation and it's requirements.